

SCATTERING IN THE PAIR OF HILBERT SPACES

E.A. Bugrij, A.V. Mishchenko¹, Yu.A. Sitenko²

Bogolyubov Institute for Theoretical Physics

252142, Kiev-142, Ukraine

Abstract

The problem of the scattering of a charged test particle in the gravitational background of axially symmetrical wormhole in the presence of the Aharonov-Bohm type magnetic field is considered. It is shown that the natural mathematical framework appropriate for the problem is the scattering theory in the pair of Hilbert spaces. Both relevant modified wave operators and S -matrices are found.

Let us consider the problem of the scattering of a charged test particle in the cylindrically-symmetric curved space with the topology of (cylinder $\times R^2$) in the presence of the static magnetic field of Aharonov-Bohm type in the sense described below. At first, we restrict ourselves by considering static cylindrically-symmetric configuration of a gravitational field with metric

$$ds^2 = dt^2 - dz^2 - f(r) [dr^2 + r^2 d\theta^2] \quad (1)$$

where r, θ are polar coordinates in the transverse z axis direction, while z is a coordinate in the direction of the space symmetry axis. For $f(r)$ only its conical behavior for $r \rightarrow \infty$ and $r \rightarrow 0$:

$$\begin{aligned} f(r) &= \left(\frac{r}{R_1}\right)^{-2\Phi_k^{(1)}}, \quad r \geq R_1, \quad \Phi_k^{(1)} < 1, \\ f(r) &= \left(\frac{r}{R_2}\right)^{-2\Phi_k^{(2)}}, \quad r \leq R_2, \quad \Phi_k^{(2)} > 1, \end{aligned} \quad (2)$$

as well as continuity condition, $f(r) \in C(0, \infty)$, are assumed. In (2), R_1 and R_2 are arbitrary constants, $0 < R_1, R_2 < \infty$. The above-mentioned gravitational structure is, in fact, a classical static wormhole, i.e. the spatial configuration consisting of two noncompact asymptotically flat conic regions connected by a compact strongly curved region – wormhole throat.

It is well known that the Aharonov-Bohm type field strength is zero in the region reachable for particles. The latter can be provided whether by puncturing the point in which the field strength is infinite (for the classical Aharonov-Bohm field) or by restriction of the particle motion to the region by means of potential barrier, for example. In our case, we can use pure geometrical device that disjoitns the region with non-zero magnetic field strength from charged matter. Namely, we can imagine that this region is concealed inside wormhole.

Motion of a test particle in the exterior along the wormhole axis is free, therefore, our concern will be on the $z = \text{const}$ section. The dynamics of the test particle under consideration is governed by the Hamiltonian:

$$H = \frac{\hbar^2}{2m} \left[-\partial_r^2 - \frac{1}{r} \partial_r + \frac{1}{r^2} \left(i\partial_\theta + \Phi + \frac{\sigma^3}{4} r \partial_r \ln f \right)^2 - \frac{1}{4} \left(\partial_r^2 \ln f + \frac{1}{r} \partial_r \ln f \right) \right], \quad (3)$$

¹ *e-mail*: olmishch@ap3.bitp.kiev.ua

² *e-mail*: yusitenko@bitp.kiev.ua

where

$$\sigma^3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix},$$

in the case of a spinor particle (ψ is a two-component column function) and $\sigma = 0$ in the case of a scalar particle (ψ is a single-component function), $0 \leq \Phi < 1$ is magnetic flux and scalar product in the Hilbert space $\mathcal{H}$ of square integrable functions on the curved space M is following:

$$(\psi, \psi') = \int_0^{2\pi} d\theta \int_0^\infty dr r f(r) \psi^\dagger \psi', \quad (4)$$

where sign $\dagger$ denotes hermitian conjugation.

It is known that to formulate the problem of scattering it is necessary, at least, to determine two Hamiltonians, H_0 and H . The first one describes non-perturbed dynamics and the second one – perturbed dynamics. Having determined the above-mentioned pair of Hamiltonians one builds wave (Möller) operators

$$\Omega^\pm(H, H_0) = s - \lim_{t \rightarrow \pm\infty} e^{iHt} e^{-iH_0 t} P(H_0). \quad (5)$$

where $P(H_0)$ is a projector on absolutely continuous subspace of operator H_0 . Let us note that expression (5) makes sense only in the case, when perturbation

$$V = H - H_0 \quad (6)$$

has a short range (see [1], [2]). For the second order operators the latter means (following Hörmander) [2] that the coefficients in the expansion

$$V = a_{ij} \partial_i \partial_j + b_i \partial_i + c \quad (7)$$

decrease like $|r|^{-1-\varepsilon}$ for $|r| \rightarrow \infty$ and $\varepsilon > 0$. For the case (following Hörmander) when perturbation has long range, when the above-mentioned (real) coefficients decrease like $|r|^{-\varepsilon}$ for $|r| \rightarrow \infty$, $\varepsilon > 0$ the definition of wave operators (5) must be modified. At last, we cannot say anything as to the existence of the wave operators in the case when Hörmander condition of long range action is broken.

We want to consider scattering of a particle which dynamics coincides with one of a free particle on the cone in the infinitely remote past. As it is known [3], the latter is described simply by free Hamiltonian on the plane R^2

$$H_0 = -\frac{\hbar^2}{2m} \Delta_2. \quad (8)$$

This is conditioned by the fact that cone is a flat surface with zero Gauss curvature everywhere except apex and, consequently, locally coincides with R^2 . Analogously, it is supposed that in the infinitely remote future reflected wave will exist. Its dynamics is determined by Hamiltonian H_0 . The wave which has passed through the wormhole throat will exist, too. And its dynamics is determined by Hamiltonian H_0 , but acting on functions that are defined on the second cone. It turns out, that in such formulation of scattering problem it is impossible to choose Hamiltonian describing a free dynamics in all curved space M .

Nevertheless, mathematical formalism appropriate for our problem exists, the scattering theory in a pair of Hilbert spaces, namely [1].

In the problem it is more convenient to use a new radial coordinate

$$\rho(r) = \int_{r_0}^r dr' [f(r')]^{1/2} \quad (9)$$

instead of r , where r_0 is arbitrary parameter. The metric (1) then has the form:

$$ds^2 = d\rho^2 + g_{00}(\rho)d\theta^2. \quad (10)$$

Let us perform the unitary mapping

$$\mathcal{H} \rightarrow \widetilde{\mathcal{H}}, \quad \psi \rightarrow \widetilde{\psi} = r^{1/2} f^{1/4} \psi \quad (11)$$

where $\widetilde{\mathcal{H}}$ is Hilbert space with scalar product

$$(\widetilde{\psi}, \widetilde{\psi}') = \int_0^{2\pi} d\theta \int_{-\infty}^{\infty} d\rho \widetilde{\psi}^\dagger \widetilde{\psi}'. \quad (12)$$

Then the Hamiltonians H and H_0 map into

$$\widetilde{H} = \frac{\hbar^2}{2m} \left[-\partial_\rho^2 + g_{00}^{-1} \left(i\partial_\theta + \phi - \frac{\sigma^3}{2} + \frac{\sigma^3}{4} g_{00}^{1/2} \partial_\rho \ln g_{00} \right)^2 - \frac{1}{16} \left(\partial_\rho \ln g_{00} \right)^2 \right], \quad (13)$$

$$\widetilde{H}_0 = -\frac{\hbar^2}{2m} \left[\partial_\rho^2 + \frac{\partial_\theta^2}{\rho^2} + \frac{1}{4\rho^2} \right]. \quad (14)$$

In the further consideration we will omit tilde sign.

In addition to the Hilbert space $\mathcal{H}$ let us define two Hilbert spaces $\mathcal{H}_+$ and $\mathcal{H}_-$. Space $\mathcal{H}_+$ is a space of square integrable functions on the right cone $\theta \in [0, 2\pi]$, $\rho \in [0, +\infty)$ with a scalar product

$$(\psi, \psi') = \int_0^{2\pi} d\theta \int_0^\infty d\rho \psi^\dagger \psi'. \quad (15)$$

Analogously, the space $\mathcal{H}_-$ is a space of square integrable functions on the left cone $\theta \in [0, 2\pi]$, $\rho \in [-\infty, 0)$ with a scalar product

$$(\psi, \psi') = \int_0^{2\pi} d\theta \int_{-\infty}^0 d\rho \psi^\dagger \psi'. \quad (16)$$

Let us also define operators $J_\pm$ which map the Hilbert spaces $\mathcal{H}_\pm$ into $\mathcal{H}$ and act as follows. For $\psi_\pm \in \mathcal{H}_\pm$:

$$J_\pm \psi_\pm = \mu_\pm(\rho) \psi_\pm, \quad (17)$$

where $\mu_\pm(\rho) \in C^1(\pm\infty, 0)$ and

$$\begin{aligned} \mu_\pm(\rho) &= 0, & |\rho| < \varepsilon, \\ \mu_\pm(\rho) &= 1, & |\rho| > \mathcal{P}, \end{aligned} \quad (18)$$

where $0 < \varepsilon < \mathcal{P} < \infty$. To solve the problem of scattering formulated earlier it is necessary to build the solution of Schrödinger equation

$$i\hbar \frac{\partial \psi}{\partial t} = H\psi, \quad (19)$$

that satisfies in the infinitely remote past the condition

$$\lim_{t \rightarrow -\infty} \|\psi(t) - J_+ e^{-iH_0^{(+)}t} \phi_+^{(-)}\| = 0 \quad (20)$$

and in the infinitely remote future the conditions

$$\begin{aligned} \lim_{t \rightarrow +\infty} \|\psi(t) - J_+ e^{-iH_0^{(+)}t} \phi_+^{(+)}\| &= 0, \\ \lim_{t \rightarrow +\infty} \|\psi(t) - J_- e^{-iH_0^{(-)}t} \phi_-^{(+)}\| &= 0. \end{aligned} \quad (21)$$

Here, $\phi_+^{(\pm)} \in \mathcal{H}_+$ and $\phi_-^{(\pm)} \in \mathcal{H}_-$, the Hamiltonians $H_0^{(+)}$ and $H_0^{(-)}$ describe a free dynamics on the right and left cones, respectively, and in compliance with (14) have the form:

$$H_0^{(\pm)} = -\frac{\hbar^2}{2m} \left[\partial_\rho^2 + \frac{\partial_0^2}{\rho^2} + \frac{1}{4\rho^2} \right]. \quad (22)$$

Two scattering matrices S_{++} and S_{+-}

$$\phi_+^{(+)} = S_{++} \phi_+^{(-)}, \quad \phi_-^{(+)} = S_{+-} \phi_+^{(-)} \quad (23)$$

will contain all necessary information about scattering. Taking into account that $\psi(t) = e^{-iHt} \psi_0$ for some $\psi_0 \in \mathcal{H}$, we get

$$\begin{aligned} S_{++} &= (\Omega_+^{(-)}(H, H_0^{(+)}; J_+))^\dagger \Omega_+^{(+)}(H, H_0^{(+)}; J_+), \\ S_{+-} &= (\Omega_-^{(-)}(H, H_0^{(-)}; J_-))^\dagger \Omega_+^{(+)}(H, H_0^{(+)}; J_+), \end{aligned} \quad (24)$$

where wave operators $\Omega_+^{(+)}$, $\Omega_+^{(-)}$, $\Omega_-^{(-)}$ are determined by expressions:

$$\begin{aligned} \Omega_+^{(+)}(H, H_0^{(+)}; J_+) &= s - \lim_{t \rightarrow -\infty} e^{iHt} J_+ e^{-iH_0^{(+)}t}, \\ \Omega_+^{(-)}(H, H_0^{(+)}; J_+) &= s - \lim_{t \rightarrow +\infty} e^{iHt} J_+ e^{-iH_0^{(+)}t}, \\ \Omega_-^{(-)}(H, H_0^{(-)}; J_-) &= s - \lim_{t \rightarrow +\infty} e^{iHt} J_- e^{-iH_0^{(-)}t}. \end{aligned} \quad (25)$$

In the expressions (25) we take into account that the spectra of operators $H_0^{(+)}$ and $H_0^{(-)}$ are absolutely continuous.

Thus, the problem of scattering is reduced to the problem of finding wave operators $\Omega_+^{(\pm)}$ for the pair of Hilbert spaces $\mathcal{H}_+$ and $\mathcal{H}$, and wave operator $\Omega_-^{(-)}$ for the pair of Hilbert spaces $\mathcal{H}_-$ and $\mathcal{H}$. Let us construct them. One can use the methods of stationary scattering theory and find wave functions $\psi_+^{(\pm)}(\vec{k})$ and $\psi_-^{(-)}(\vec{k})$ instead of wave operators. These functions are the result of action of wave operators on functions

$$\psi_\pm^{(0)}(k) = \frac{1}{2\pi} e^{ik\rho \cos(\theta - \theta_k)}, \quad (26)$$

where θ_k is the angle that determines the direction of wave vector $\vec{k}$ and $k = |\vec{k}|$; $\rho \in (-\infty, 0)$ for $\psi_-^{(0)}$ and $\rho \in (0, +\infty)$ for $\psi_+^{(0)}$ and the functions $\psi_\pm^{(0)}$ are free stationary waves on the right and left cones. Then

$$\psi_+^{(\pm)}(\vec{k}) = \Omega_+^{(\pm)} \psi_\pm^{(0)}(\vec{k}), \quad \psi_-^{(\pm)}(\vec{k}) = \Omega_-^{(-)} \psi_-^{(0)}(\vec{k}). \quad (27)$$

While expansion on functions $\psi_{\pm}^{(0)}(\vec{k})$ is, in fact, Furier transformation for spaces $\mathcal{H}_{\pm}$, then functions $\psi_{+}^{(\pm)}(\vec{k})$ and $\psi_{-}^{(\pm)}(\vec{k})$ completely define the wave operators $\Omega_{+}^{(\pm)}$ and $\Omega_{-}^{(\pm)}$. Using expressions (25) and (26), (27), and the fact that

$$H_0^{(\pm)}\psi_{\pm}^{(0)} = \frac{\hbar^2 k^2}{2m} \quad (28)$$

as well, one can simply get the following Lippmann-Shwinger equations:

$$\begin{aligned} \psi_{+}^{(\pm)}(\vec{k}) &= J_{+}\psi_{+}^{(0)}(\vec{k}) - G^{\pm}(\vec{k})[HJ_{+} - J_{+}H_0^{(+)}]\psi_{+}^{(0)}(\vec{k}), \\ \psi_{-}^{(\pm)}(\vec{k}) &= J_{-}\psi_{-}^{(0)}(\vec{k}) - G^{\pm}(\vec{k})[HJ_{-} - J_{-}H_0^{(-)}]\psi_{-}^{(0)}(\vec{k}), \end{aligned} \quad (29)$$

where Green functions

$$G^{\pm}(\vec{k}) = \lim_{\varepsilon \rightarrow +0} \left[H - \frac{\hbar^2 k^2}{2m} \mp i\varepsilon \right]^{-1}. \quad (30)$$

Taking into account the cylindrical symmetry of the problem, let us expand the functions $\psi_{+}^{(\pm)}$ and $\psi_{-}^{(\pm)}$ in the Furier series:

$$\psi_{+}^{(\pm)} = \sum_{n=-\infty}^{+\infty} e^{in\theta} \psi_{+,n}^{(\pm)}, \quad \psi_{-}^{(\pm)} = \sum_{n=-\infty}^{+\infty} e^{in\theta} \psi_{-,n}^{(\pm)}. \quad (31)$$

The Hilbert space $\mathcal{H}$ will expand, correspondingly, to the direct sum $\mathcal{H} = \bigoplus_{n=-\infty}^{n=+\infty} \mathcal{H}_n$, and scalar product on $\mathcal{H}_n$ is defined as follows:

$$(\psi_n, \psi'_n) = \int_{-\infty}^{+\infty} d\rho \psi_n^{\dagger} \psi'_n.$$

To construct the Green functions it is necessary to calculate the resolvent of operator H :

$$R_z = (H - z)^{-1}. \quad (32)$$

Let us choose the branch of $\sqrt{z}$ that $\Im \sqrt{z} \geq 0$ and fix that $\omega = \frac{\hbar}{\sqrt{2m}} \sqrt{z}$. Then it is easy to get the following expression for the kernel of the revolvent of operator H :

$$R_z(\rho, \rho', \theta, \theta') = \frac{1}{2\pi} \sum_{n=-\infty}^{+\infty} e^{in(\theta-\theta')} R_n(\rho, \rho'), \quad (33)$$

where

$$R_n(\rho, \rho') = \frac{i2m}{2a_n(\omega)\omega\hbar^2} \left[\psi_1^{(n)}(\rho, \omega) \psi_2^{(n)}(\rho', \omega) \theta(\rho - \rho') + \psi_2^{(n)}(\rho, \omega) \psi_1^{(n)}(\rho', \omega) \theta(\rho' - \rho) \right]. \quad (34)$$

In expression (34)

$$\theta(x) = \begin{cases} 1, & x > 0 \\ 0, & x < 0 \end{cases}$$

and

$$a_n(\omega) = \frac{i}{2\omega} W\{\psi_1^{(n)}, \psi_2^{(n)}\},$$

where for two functions $f_1(\rho)$ and $f_2(\rho)$ Wronskian is defined as follows

$$W\{f_1(\rho), f_2(\rho)\} = f_1(\rho)\partial_\rho f_2(\rho) - f_2(\rho)\partial_\rho f_1(\rho).$$

The functions $\psi_1^{(n)}(\rho, \omega)$ and $\psi_2^{(n)}(\rho, \omega)$ are the pair of linearly independent Jost solutions for partial Hamiltonian $H_n = e^{-in\theta} H e^{in\theta}$, which are characterized by following asymptotics:

$$\begin{aligned}\psi_1^{(n)}(\rho, \omega) &= e^{i\omega\rho}, & \rho \rightarrow +\infty, \\ \psi_2^{(n)}(\rho, \omega) &= e^{-i\omega\rho}, & \rho \rightarrow -\infty.\end{aligned}\tag{35}$$

Having defined the second pair of linearly independent Jost solutions

$$\begin{aligned}\bar{\psi}_1^{(n)}(\rho, \omega) &= e^{-i\omega\rho}, & \rho \rightarrow +\infty, \\ \bar{\psi}_2^{(n)}(\rho, \omega) &= e^{i\omega\rho}, & \rho \rightarrow -\infty,\end{aligned}\tag{36}$$

one can write the following useful expressions:

$$\psi_1^{(n)} = a_n(\omega)\bar{\psi}_2^{(n)} + b_n(\omega)\psi_2^{(n)}, \quad \psi_2^{(n)} = a_n(\omega)\bar{\psi}_1^{(n)} - \bar{b}_n(\omega)\psi_1^{(n)};\tag{37}$$

$$\bar{b}_n(\omega) = (b_n(\omega))^* = b_n(-\omega^*), \quad (a_n(\omega))^* = a_n(-\omega^*);\tag{38}$$

$$\bar{\psi}_i^{(n)}(\rho, \omega) = \psi_i^{(n)}(\rho, -\omega) = (\psi_i^{(n)}(\rho, \omega^*))^*, \quad i = 1, 2;\tag{39}$$

$$\begin{aligned}W\{\psi_1^{(n)}(\rho, \omega), \bar{\psi}_2^{(n)}(\rho, \omega)\} &= 2ib_n(\omega)\omega, \\ W\{\psi_1^{(n)}(\rho, \omega), \psi_2^{(n)}(\rho, \omega)\} &= -2ia_n(\omega)\omega, \\ W\{\psi_2^{(n)}(\rho, \omega), \bar{\psi}_2^{(n)}(\rho, \omega)\} &= -W\{\psi_1^{(n)}(\rho, \omega), \bar{\psi}_1^{(n)}(\rho, \omega)\} = 2i\omega.\end{aligned}\tag{40}$$

For real ω let us fix $k = \omega$.

Then the following expression takes place:

$$|a_n(k)|^2 - |b_n(k)|^2 = 1.\tag{41}$$

For the partial reflection and transmission coefficients one can write

$$r_n(k) = b_n(k)/a_n(k), \quad t_n(k) = 1/a_n(k).\tag{42}$$

Then

$$|r_n(k)|^2 + |t_n(k)|^2 = 1\tag{43}$$

is partial unitarity condition.

The following scalar products

$$\begin{aligned}(\psi_1^{(n)}(k), \psi_1^{(n)}(k')) &= 2\pi|a_n(k)|^2\delta(k - k'), \\ (\psi_2^{(n)}(k), \psi_2^{(n)}(k')) &= 2\pi|a_n(k)|^2\delta(k - k'), \quad k, k' > 0, \\ (\psi_1^{(n)}(k), \psi_2^{(n)}(k')) &= 0,\end{aligned}\tag{44}$$

are necessary for the further consideration.

Expanding Green functions

$$G_n^+(\rho, \rho') = \frac{1}{2\pi} \sum_{n=-\infty}^{\infty} e^{in(\theta-\theta')} G_n^\pm(\rho, \rho') \quad (45)$$

and using expressions (30), (34), (38), (39) one can find for partial Green functions

$$\begin{aligned} G_n^+(\rho, \rho') &= \frac{im}{a(k)k\hbar^2} [\psi_1^{(n)}(\rho, k) \psi_2^{(n)}(\rho', k') \theta(\rho - \rho') + \\ &\quad + \psi_2^{(n)}(\rho, k) \psi_1^{(n)}(\rho', k) \theta(\rho' - \rho)], \\ G_n^-(\rho, \rho') &= (G_n^+(\rho, \rho'))^* \end{aligned} \quad (46)$$

Substituting (31), (45), (46) in the Lippmann-Schwinger equations of (29) and taking into account the expansion

$$\psi_\pm^{(0)}(k) = \frac{e^{ik\rho \cos(\theta-\theta_k)}}{2\pi} = \frac{\sqrt{\rho}}{2\pi} \sum_{n=-\infty}^{+\infty} e^{in(\theta-\theta_k) + \frac{i\pi|n|}{2}} J_{|n|}(k\rho), \quad (47)$$

we get:

$$\psi_{+,n}^{(+)} = \frac{\psi_2^{(n)}}{2\pi a_n(k)} \sqrt{\frac{1}{2\pi k}} e^{-in\theta_k + i\pi|n| + \frac{i\pi}{4}}, \quad (48)$$

$$\psi_{+,n}^{(-)} = \frac{(\psi_2^{(n)})^*}{2\pi a_n^*(k)} \sqrt{\frac{1}{2\pi k}} e^{-in\theta_k - \frac{i\pi}{4}}, \quad (49)$$

$$\psi_{-,n}^{(-)} = \frac{(\psi_1^{(n)})^*}{2\pi a_n^*(k)} \sqrt{\frac{1}{2\pi k}} e^{-in\theta_k - \frac{i\pi}{4}}. \quad (50)$$

Let us find, now, operators S_{++} and S_{+-} . They are completely defined by following matrix elements:

$$\begin{aligned} S_{++}(\vec{k}, \vec{k}') &= (\psi_+^{(0)}(\vec{k}), S_{++}\psi_+^{(0)}(\vec{k}'))_{\mathcal{H}_+} = (\psi_+^{(-)}(\vec{k}), \psi_+^{(+)}(\vec{k}'))_{\mathcal{H}} \\ S_{+-}(\vec{k}, \vec{k}') &= (\psi_+^{(0)}(\vec{k}), S_{+-}\psi_-^{(0)}(\vec{k}'))_{\mathcal{H}_+} = (\psi_-^{(-)}(\vec{k}), \psi_+^{(+)}(\vec{k}'))_{\mathcal{H}} \end{aligned} \quad (51)$$

Using expressions (31), (44), (48)–(50) one can find for matrix elements (51):

$$S_{++}(\vec{k}, \vec{k}') = -\frac{1}{2\pi\sqrt{k k'}} e^{i\pi/2} \delta(k - k') \sum_{n=-\infty}^{+\infty} e^{in(\theta_k - \theta_{k'}) + i\pi|n|} \frac{b_n^*(k)}{a_n(k)}, \quad (52)$$

$$S_{+-}(\vec{k}, \vec{k}') = \frac{1}{2\pi\sqrt{k k'}} e^{i\pi/2} \delta(k - k') \sum_{n=-\infty}^{+\infty} e^{in(\theta_k - \theta_{k'}) + i\pi|n|} \frac{1}{a_n(k)}. \quad (53)$$

The kernel of operator S_{++} (52) can be written in terms of the sum of singular and regular, with respect to the angular variable, functions. Actually, correspondingly to (2), (9), (13) and (35)

$$\begin{aligned} \psi_1^{(n)} &= \sqrt{\frac{\pi k(\rho + \rho_1)}{2}} e^{i(-k\rho_1 + \frac{\pi\alpha_n^{(1)}}{2} + \frac{\pi}{4})} H_{\alpha_n^{(1)}}^{(1)}(k(\rho + \rho_1)), \quad \rho \geq P_1, \\ \psi_2^{(n)} &= \sqrt{\frac{\pi k(\rho + \rho_2)}{2}} e^{i(k\rho_2 + \frac{\pi\alpha_n^{(2)}}{2} + \frac{\pi}{4})} H_{\alpha_n^{(2)}}^{(1)}(k|\rho + \rho_2|), \quad \rho \leq P_2 \end{aligned} \quad (54)$$

Using (39) we get from (54)

$$\begin{aligned}\bar{\psi}_1^{(n)} &= \sqrt{\frac{\pi k(\rho + \rho_1)}{2}} e^{i(k\rho_1 - \frac{\pi\alpha_n^{(1)}}{2} - \frac{\pi}{4})} H_{\alpha_n^{(1)}}^{(2)}(k(\rho + \rho_1)), \\ \bar{\psi}_2^{(n)} &= \sqrt{\frac{\pi k(\rho + \rho_2)}{2}} e^{i(-k\rho_2 - \frac{\pi\alpha_n^{(2)}}{2} - \frac{\pi}{4})} H_{\alpha_n^{(2)}}^{(2)}(k(\rho + \rho_2)).\end{aligned}\quad (55)$$

where

$$\begin{aligned}P_i &= \int_{r_0}^{R_i} dr' \sqrt{f(r')}, \\ \alpha_n^{(i)} &= \left| \frac{n - \Phi + \frac{1}{2}\sigma^3 \Phi_k^{(i)}}{1 - \Phi_k^{(i)}} \right|, \quad i = 1, 2, \\ \rho_i &= \frac{R_i}{1 - \Phi_n^{(i)}} - P_i.\end{aligned}\quad (56)$$

From (40) we find then:

$$\frac{b_n^*(k)}{a_n(k)} = \frac{W(\bar{\psi}_n^{(1)}(k), \psi_n^{(2)}(k))}{W(\psi_n^{(1)}(k), \psi_n^{(2)}(k))}. \quad (57)$$

As is known, for large order and finite value of argument of Bessel functions

$$H_\nu^{(1)}(z) = -H_\nu^{(2)}(z) \left[1 + O\left(\frac{ez}{2\nu}\right)^{2\nu} \right]. \quad (58)$$

Then, it follows from (54), (55), (57), (58) that

$$b_n^*(k) = -e^{2i(k\rho_1 - \frac{\pi\alpha_n^{(1)}}{2} - \frac{\pi}{4})} a_n(k) \left\{ 1 + O\left[\left(\frac{ek(P_1 + \rho_1)}{2\alpha_n^{(1)}}\right)^{2\alpha_n^{(1)}}\right] \right\} \quad (59)$$

Using partial expression of unitarity (41) we find that

$$\frac{1}{|a_n|^2} = 1 - \frac{|b_n|^2}{|a_n|^2} = 1 - \left| 1 + O\left[\left(\frac{ek(P_1 + \rho_1)}{2\alpha_n^{(1)}}\right)^{2\alpha_n^{(1)}}\right] \right|^2 = O\left(\frac{ek(P_1 + \rho_1)}{2\alpha_n^{(1)}}\right)^{2\alpha_n^{(1)}}. \quad (60)$$

For small values of argument $(k(P_1 + \rho_1))$ and $\alpha_n^{(1)} \neq 0$ the following expressions take place:

$$H_{\alpha_n^{(1)}}^{(2)}(k(P_1 + \rho_1)) = -H_{\alpha_n^{(1)}}^{(1)}[1 + O[(k(P_1 + \rho_1))^{\alpha_n^{(1)}}]] \quad (61)$$

and for $\alpha_n^{(1)} = 0$:

$$H_0^{(2)}(k(P_1 + \rho_1)) = -H_0^{(1)}(k(P_1 + \rho_1)) \left[1 + O\left(\frac{1}{|\ln(k(P_1 + \rho_1))|}\right) \right], \quad (62)$$

$$\begin{aligned}b_n^*(k) &= -e^{2i(k\rho_1 - \frac{\pi\alpha_n^{(1)}}{2} - \frac{\pi}{4})} a_n(k) \left\{ 1 + O\left[\left(k(P_1 + \rho_1)\right)^{\alpha_n^{(1)}}\right] \right\}, \quad \alpha_n^{(1)} \neq 0, \\ b_n^*(k) &= -e^{2i(k\rho_1 - \frac{\pi\alpha_n^{(1)}}{2} - \frac{\pi}{4})} a_n(k) \left\{ 1 + O\left[\left(\frac{1}{|\ln(k(P_1 + \rho_1))|}\right)\right] \right\}, \quad \alpha_n^{(1)} = 0.\end{aligned}\quad (63)$$

$$\begin{aligned}\frac{1}{|a_n|^2} &= 1 - \frac{|b_n|^2}{|a_n|^2} = O\left[\left(k(P_1 + \rho_1)\right)^{\alpha_n^{(1)}}\right], \quad \alpha_n^{(1)} \neq 0, \\ \frac{1}{|a_n|^2} &= 1 - \frac{|b_n|^2}{|a_n|^2} = O\left[\left(\frac{1}{|\ln(k(P_1 + \rho_1))|}\right)\right], \quad \alpha_n^{(1)} = 0.\end{aligned}\quad (64)$$

It follows from (60) that series (53) for the kernel of the operator $S_{+-}(\vec{k}, \vec{k}')$ converges uniformly with respect to the angular variable and is therefore regular function of the angular variable. Moreover, this function is infinitely differentiable with respect to the angular variable and, in accordance with (64), behaves like $O\left[(k(P_1 + \rho_1))^{\alpha_n^{(1)}-1}\right]$ or $O\left[(k|\ln(k(P_1 + \rho_1))|)^{-1}\right]$ when $k(P_1 + \rho_1) \rightarrow 0$.

Using (59) and (63) one can represent the kernel of the operator $S_{++}(\vec{k}, \vec{k}')$ (52) in the following form:

$$\begin{aligned}S_{++}(\vec{k}, \vec{k}') &= \frac{\delta(k - k')}{2\pi\sqrt{kk'}} e^{2ik\rho_1} \sum_{n=-\infty}^{\infty} e^{in(\theta_k - \theta_{k'}) + i\pi|n| - i\pi\alpha_n^{(1)}} - \\ &- \frac{\delta(k - k')}{2\pi\sqrt{kk'}} e^{2ik\rho_1} \sum_{n=-\infty}^{\infty} e^{in(\theta_k - \theta_{k'}) + i\pi|n| - i\pi\alpha_n^{(1)}} \times \\ &\times \left[\frac{b_n^*(k) e^{i\pi/2 + i\pi\alpha_n^{(1)} - 2ik\rho_1} + a_n(k)}{a_n(k)} \right].\end{aligned}\quad (65)$$

The first sum in (65) is the kernel of S -matrix that describes scattering of particle on coinciding singular gravitational and magnetic strings. They are characterized by the curvature flux $\Phi_K^{(1)}$ and magnetic flux Φ respectively. We can write this sum as follows

$$I'(k, \theta_k; k', \theta_{k'}) + \delta(k - k') \sqrt{\frac{i}{2\pi k}} f(k; \theta_k - \theta_{k'}) e^{2ik\rho_1}, \quad (66)$$

where I' is modified unity operator. It is characterized by singular kernel

$$\begin{aligned}I'(k, \theta_k; k', \theta_{k'}) &= \frac{1}{2} \frac{\delta(k - k')}{\sqrt{kk'}} e^{2ik\rho_1} \times \\ &\times \left\{ \Delta(\theta_k - \theta_{k'} + \beta_1\pi) \exp[-i\mu_1(1 + \beta_1)\pi] + \Delta(\theta_k - \theta_{k'} - \beta_1\pi) \exp[i\mu_1(1 + \beta_1)\pi] \right\}\end{aligned}\quad (67)$$

where $\Delta(\theta) = \frac{1}{2\pi} \sum_{n=-\infty}^{\infty} e^{in\theta}$ – periodic delta-function, $\mu_1 = \Phi - \sigma^3 \Phi_K^{(1)}/2$; $\beta_1 = \Phi_K^{(1)}(1 - \Phi_K^{(1)})^{-1}$; f – scattering amplitude of particle on coinciding singular gravitational and magnetic strings.

The second sum in (65) is regular infinitely differentiable function with respect to the angular variable and its behavior is the same as for $S_{+-}(\vec{k}, \vec{k}')$ for $k(P_1 + \rho_1) \rightarrow 0$. One can also formulate the problem of scattering of particle directed from “right” to “left” cone. For solving this problem it is necessary to define additionally the operator

$$\Omega_-^{(+)}(H, H_0^{(-)}; J_-) = s - \lim_{t \rightarrow -\infty} e^{iHt} J_- e^{-iH_0^{(-)}t} \quad (68)$$

or in terms of stationary scattering theory, wave function

$$\psi_-^{(+)} = \Omega_-^{(+)}(H, H_0^{(-)}; J_-) \psi_-^{(0)}. \quad (69)$$

It is easy to find it by the way same as above

$$\psi_{-,n}^{(+)} = \frac{\psi_n^{(1)}}{2a_n(k)} \sqrt{\frac{2}{\pi k}} e^{-in\theta_k + i\pi/4 + i\pi|n|}. \quad (70)$$

In addition to the matrices of scattering S_{++} and S_{+-} that describe probability of reflection to “left” cone and transmission to the “right” cone, respectively, for particle directed from left to right, we have to define matrices S_{--} and S_{-+} describing reflection to the “right” cone and transmission to “left” one:

$$\begin{aligned} S_{--} &= (\Omega_-^{(-)}(H, H_-^{(0)}; J_-))^\dagger \Omega_-^{(+)}(H, H_-^{(0)}; J_-); \\ S_{-+} &= (\Omega_+^{(-)}(H, H_+^{(0)}; J_-))^\dagger \Omega_-^{(+)}(H, H_-^{(0)}; J_-); \end{aligned} \quad (71)$$

They are completely determined by their matrix elements:

$$\begin{aligned} S_{--}(\vec{k}, \vec{k}') &= (\psi_-^{(0)}(\vec{k}), S_{--}\psi_-^{(0)}(\vec{k}'))_{\mathcal{H}_-} = (\psi_-^{(-)}(\vec{k}), \psi_-^{(+)}(\vec{k}'))_{\mathcal{H}}, \\ S_{-+}(\vec{k}, \vec{k}') &= (\psi_-^{(0)}(\vec{k}), S_{-+}\psi_+^{(0)}(\vec{k}'))_{\mathcal{H}_-} = (\psi_+^{(-)}(\vec{k}), \psi_-^{(+)}(\vec{k}'))_{\mathcal{H}}. \end{aligned} \quad (72)$$

Finally:

$$\begin{aligned} S_{--}(\vec{k}, \vec{k}') &= \frac{\delta(k - k')}{2\pi\sqrt{kk'}} e^{i\pi/2} \sum_{n=-\infty}^{\infty} e^{in(\theta_k - \theta_{k'}) + i\pi|n|} \frac{b_n(k)}{a_n(k)}, \\ S_{-+}(\vec{k}, \vec{k}') &= S_{+-}(\vec{k}, \vec{k}') \end{aligned} \quad (73)$$

It is possible to write the kernel of operator $S_{--}(\vec{k}, \vec{k}')$ in terms of the sum of singular and regular functions with respect to the angular variable:

$$\begin{aligned} S_{--} &= \frac{\delta(k - k')}{2\pi\sqrt{kk'}} \sum_{n=-\infty}^{\infty} e^{in(\theta_k - \theta_{k'}) + i\pi|n| - i\pi\alpha_n^{(2)} - 2ik\rho} + \\ &+ \frac{\delta(k - k')}{2\pi\sqrt{kk'}} \sum_{n=-\infty}^{\infty} e^{in(\theta_k - \theta_{k'}) + i\pi|n| - i\pi\alpha_n^{(2)} - 2ik\rho_2} \times \\ &\times \left[\frac{b_n(k) e^{i\pi\alpha_n^{(2)} + i\pi/2 + 2ik\rho_2} - a_n(k)}{a_n(k)} \right]. \end{aligned} \quad (74)$$

From expressions (52)–(53), using the partial expression of unitarity (41) it is easy to get the following expression

$$(S_{++}S_{++}^\dagger)(\vec{k}, \vec{k}') + (S_{+-}S_{+-}^\dagger)(\vec{k}, \vec{k}') = \frac{\delta(k - k')}{\sqrt{kk'}} \Delta(\theta_k - \theta_{k'}). \quad (75)$$

Analogously for S_{--} , S_{-+}

$$(S_{--}S_{--}^\dagger)(\vec{k}, \vec{k}') + (S_{-+}S_{-+}^\dagger)(\vec{k}, \vec{k}') = \frac{\delta(k - k')}{\sqrt{kk'}} \Delta(\theta_k - \theta_{k'}). \quad (76)$$

The equations (75)–(76) are unitarity conditions for the problem under consideration.

Acknowledgments

The work of A.V.M. and Yu.A.S. was supported by Swiss National Science Foundation Grant No. CEEC/NIS/96-98/7 IP 051219.

References

- [1] M.C. Reed and B. Simon, *Method of Modern Mathematical Physics*, Vol. 3. Scattering Theory, 1979.
- [2] L. Hörmander, *Analysis of Linear Partial Differential Operators*, II, IV, 1985.
- [3] Yu.A.Sitenko, A.V.Mishchenko, *Aharonov-Bohm effect in curved space and cosmic strings*, Zh.Eksp.Teor. Fiz., vol. 1-8, 1516–1553, 1995.